



GASTROINTESTINAL SYMPTOMS IN ENTERALLY FED ADULTS WITH NEUROLOGICAL CONDITIONS

A multidisciplinary tool to aid management


Developed by Anton Emmanuel (Consultant Neuro-Gastroenterologist, The National Hospital for Neurology and Neurosurgery, London), Tanita Manton (Dietitian, The Royal Hospital for Neuro-disability, London), Emma Carder (Dietitian, The Walton Centre, Liverpool), and Carolyn Street (Dietitian, Regional Hyper-acute Rehabilitation Unit, Northwick Park Hospital, London), with the support of Nestlé Health Science UK.

This multidisciplinary tool is experience-based, and where insufficient evidence exists, recommendations are based on consensus opinions of best practice. Each patient is an individual – this tool has been developed to support clinical practice and should be used in conjunction with clinical judgement and patient wishes. Nestlé Health Science did not influence decision-making during the development of this resource – the views and recommendations expressed are those of the authors.



A RECENT UK AUDIT SHOWED THAT 28% OF LONG-TERM PATIENTS WITH NEUROLOGICAL DISORDERS HAD DOCUMENTED EVIDENCE OF ENTERAL FEEDING INTOLERANCE¹

While GI dysfunction and enteral feeding intolerance is well documented in children with neurological disorders² and in other adult patient groups,³ evidence is comparatively sparse regarding adult neurological disorders.





A RECENT UK AUDIT SHOWED THAT 28% OF LONG-TERM PATIENTS WITH NEUROLOGICAL DISORDERS HAD DOCUMENTED EVIDENCE OF ENTERAL FEEDING INTOLERANCE.¹

INTRODUCTION

ASSESSMENT AND
PATIENT HISTORY

UPPER
GI SYMPTOMS

LOWER
GI SYMPTOMS

CONCLUSION

Although there is currently no consensus definition of 'enteral feeding intolerance',^{4,5} clinical manifestations may include nausea, vomiting or regurgitation, diarrhoea, abdominal pain, abdominal distension and high gastric residual volumes.³⁻⁵ These symptoms may hinder the delivery of enteral formula, with implications for nutritional status. The presence of a neurological disorder may further complicate this clinical picture, owing to the complex gut-brain relationship and medical treatments that may impact on GI function.

To help guide clinical practice in this complex area, a working group of specialist dietitians and a consultant neuro-gastroenterologist met in January 2018 with the aim of developing a pragmatic guidance tool. Clinical experience and relevant evidence were drawn upon to form consensus opinions. The need for pragmatic guidance based on clinical judgement and best practice was highlighted where there was a lack of supporting data. Design of the tool was based on the discussions of the working group, the details of which are documented here.





A RECENT UK AUDIT SHOWED THAT 28% OF LONG-TERM PATIENTS WITH NEUROLOGICAL DISORDERS HAD DOCUMENTED EVIDENCE OF ENTERAL FEEDING INTOLERANCE.¹

INTRODUCTION

ASSESSMENT AND
PATIENT HISTORY

UPPER
GI SYMPTOMS

LOWER
GI SYMPTOMS

CONCLUSION

Taking a structured patient history may help to both identify potential causative factors for GI symptoms and determine their severity. The first stage is to establish what was 'normal' for the patient prior to symptom onset. It may be possible to identify past dietary triggers to help optimise current nutritional management (e.g. modulation of fibre intake). A history of recent medication changes may help identify potential iatrogenic cause. Alarm features ('red flags') may help identify organic disease, such as ischaemic colitis, poor liver function, intracranial causes of vomiting, IBD and very rarely GI auras for epilepsy or 'abdominal' epilepsy.

The use of assessment tools can be particularly helpful in establishing the severity of current symptoms. The King's Stool Chart, which is validated in enterally-fed populations,^{6,7} may help to more accurately quantify stool volume and type. A 'vomiting diary' that records the details of, and circumstances surrounding, any vomiting episodes may highlight the severity of upper GI symptoms and elucidate potential triggers. In the critical care setting, gastric residual volumes may also be recorded.





A RECENT UK AUDIT SHOWED THAT 28% OF LONG-TERM PATIENTS WITH NEUROLOGICAL DISORDERS HAD DOCUMENTED EVIDENCE OF ENTERAL FEEDING INTOLERANCE.¹

INTRODUCTION

ASSESSMENT AND PATIENT HISTORY

UPPER GI SYMPTOMS

LOWER GI SYMPTOMS

CONCLUSION

Upper GI symptoms may include reflux, vomiting, epigastric pain, bloating and early satiety. Due to the communication difficulties that some patients experience, identification of less obvious symptoms such as reflux and indigestion may be more challenging. Patients may report or signal that they are experiencing abdominal pain. It would help to differentiate between pain localised to the stomach (**epigastric pain**), which may be related to indigestion, and the pain of acid **reflux** into the oesophagus with pain behind the sternum, in the throat or chest. In patients with asymptomatic reflux, surrogate markers may include chest infections, nocturnal cough, brash and low pH aspirates from suction aids of tracheostomies.

Nausea, vomiting and **early satiety** (feeling of fullness) after feeding may be caused by delayed gastric emptying. A 'vomiting diary' may identify triggers such as manual handling, tracheostomy suctioning, administration of medications or large boluses of fluid via the feeding tube. While it is important to consider non-feed-related factors, helpful nutrition-related strategies including reducing the size of fluid boluses or using a more concentrated enteral feed can be tried proactively to help alleviate symptoms.

Where first-line strategies fail, a change of formula may be appropriate; for example, whey-based peptide formulas may help to enhance gastric emptying.^{8,9} The exclusion of constipation as a contributing factor is also important. Medications that may be helpful for upper GI symptoms include proton pump inhibitors and prokinetics such as domperidone; however, metoclopramide should be avoided in Parkinson's disease and traumatic brain injury.¹⁰ Where vomiting episodes are linked to meals, a peripherally acting anti-emetic such as domperidone or metoclopramide may be effective.

Another common phenomenon is the sensation of **hunger** (often despite meeting full nutritional requirements with enteral formula). This can be worrying for families. In some cases, however, the patient may be expressing a desire to eat, rather than true physical hunger. Careful weight monitoring may provide reassurance that the patient is receiving adequate nutrition.

Provision of calories should not be increased instinctively, as patients with low mobility are particularly at risk of unwanted weight gain. If bolus feeding is feasible, this may help mimic a normal meal pattern. For patients who are nil by mouth and wish to eat, the sight and smell of food may be particularly difficult. If possible, food service should be handled discretely.





A RECENT UK AUDIT SHOWED THAT 28% OF LONG-TERM PATIENTS WITH NEUROLOGICAL DISORDERS HAD DOCUMENTED EVIDENCE OF ENTERAL FEEDING INTOLERANCE.¹

INTRODUCTION

ASSESSMENT AND PATIENT HISTORY

UPPER GI SYMPTOMS

LOWER GI SYMPTOMS

CONCLUSION

Constipation, diarrhoea, abdominal pain and bloating may all complicate enteral feeding. As with upper GI symptoms, establishing the likely cause by taking a detailed patient history is important. It is worthwhile asking the patient or their relatives if there is a pre-morbid history of bowel problems such as IBS or constipation.

In **constipation**, it is important to establish whether this is a result of slow transit, or rectal evacuation difficulties. Slow transit is hallmarked by infrequent urge and hard pellet stools, while the symptoms of evacuation disorder comprise need to strain, a sense of incomplete emptying or having to digitally assist voiding. For the latter, MASCIP guidelines¹¹ provide helpful information on management. In cases of slow transit, senna, stool softeners and prokinetics may all be useful, although lactulose should be avoided as it can worsen symptoms of pain and bloating.¹²

Based on a paediatric study and personal experience within the working group, in patients with persistent constipation it may be beneficial to trial a whey-based peptide formula to help relieve symptoms of pain and bloating and improve tolerance to enteral feed.¹³

Diarrhoea is defined by the World Health Organization as 'the passage of three or more loose or liquid stools per day (or more frequent passage than is normal for the individual)'.¹⁴ In an acute patient, diarrhoea should trigger onward referral, and it is important to rule out constipation and infective causes. Certain medications commonly used in neurological disorders may worsen diarrhoea, for example baclofen, antibiotics or preparations containing sorbitol.¹⁵ Anti-diarrhoeals may be explored with the medical team, and loperamide syrup may be particularly suitable as the dose can be more sensitively titrated according to the patient's changing needs.

For both constipation and diarrhoea, modulating the amount and type of fibre provided by enteral feeding can be an effective strategy. Fibre intakes of 20–25g/day for women and 30–35g/day for men are recommended for healthy adults.¹⁶

Several studies have highlighted the benefits of fibre for reducing the risk of diarrhoea in enterally-fed patients.^{17,18} Healthcare professionals may choose between different sources of fibre (soluble, insoluble, or mixed), although more evidence is needed to inform the selection or avoidance of these. The addition of fibre supplements to non-fibre containing formulas may provide a flexible approach and allow for more sensitive adjustment of doses. The benefits of partially hydrolysed guar gum (PHGG, a soluble fibre) for reducing diarrhoea have been highlighted in several publications,^{19–21} while evidence also supports its role in the management of constipation.²⁰

For some patients, however, fibre may exacerbate symptoms and therefore the reduction or avoidance of fibre may be appropriate.²¹ In addition, formulas that are lower in FODMAPs (fermentable oligosaccharides, disaccharides, monosaccharides and polyols), e.g. fructo-oligosaccharides (FOS) and inulin, have also been associated with a lower risk of, and improvements in, diarrhoea.^{22,23}

Bloating may also respond to a change in the amount or type of fibre given, or venting of air from the feeding tube using a syringe. Any adjustment to fibre intake should be gradual, and the effect on GI function monitored. Peptide based enteral formulas may also be utilised for managing diarrhoea where other strategies have not provided adequate relief.²⁴

The role of probiotics was discussed during the development of this resource, but it was felt the available evidence is too limited in this patient group to reach a consensus.





A RECENT UK AUDIT SHOWED THAT 28% OF LONG-TERM PATIENTS WITH NEUROLOGICAL DISORDERS HAD DOCUMENTED EVIDENCE OF ENTERAL FEEDING INTOLERANCE.¹

INTRODUCTION

ASSESSMENT AND
PATIENT HISTORY

UPPER
GI SYMPTOMS

LOWER
GI SYMPTOMS

CONCLUSION

GI dysfunction in this patient group may be a common, yet under-recognised problem, therefore MDT collaboration is essential in its management. This pragmatic guidance tool may be used in conjunction with clinical judgement, patient wishes and wider MDT input. It may be beneficial to trial one management strategy at a time to identify which intervention has improved symptoms.

In practice, however, it may be appropriate to consider medical and nutritional interventions concurrently. Ongoing review of the literature to support this document is planned and it is hoped that this tool will be used and audited to identify if there is a contribution to patient care.




References

1. Parry S. *ESPEN Congress*. 2016; Poster MON-P098.
2. Quitadamo P, et al. *Eur J Paediatr Neurol*. 2016; 20(6): 810–15.
3. Blaser AR, et al. *Acta Anaesthesiol Scand*. 2014; 58(8): 914–22.
4. Blaser AR, et al. *Clin Nutr*. 2015; 34(5): 956–61.
5. Wang K, et al. *JPEN J Parenter Enteral Nutr*. 2017; 41(6): 959–67.
6. Whelan K, et al. *Eur J Clin Nutr*. 2004; 58(7): 1030–7.
7. King's College London. King's Stool Chart. Available at: www.kcl.ac.uk/stoolchart. Accessed March 2018.
8. Romano C, et al. *J Pediatr Gastroenterol Nutr*. 2017; 65(2): 242–264.
9. Alexander DD, et al. *World J Gastrointest Pharmacol Ther*. 2016; 7(2): 306–19.
10. Shin HW, Chung S. *J Clin Neurol*. 2012; 8(1): 15–21.
11. Multidisciplinary Association of Spinal Cord Injured Professionals. *Guidelines for Management of Neurogenic Bowel Dysfunction in Individuals with Central Neurological Conditions*. MASCIP, 2012. Available at: <https://www.mascip.co.uk/wp-content/uploads/2015/02/CV653NNeurogenic-Guidelines-Sept-2012.pdf>. Accessed March 2018.
12. Lee-Robichaud H, et al. *Cochrane Database Syst Rev*. 2010; 7: CD007570.
13. Minor G, et al. *Glob Pediatr Health*. 2016; 21; 3:2333794X16681887.
14. World Health Organization. Diarrhoeal disease. WHO, 2017. Available at: <http://www.who.int/mediacentre/factsheets/fs330/en/>. Accessed March 2018.
15. Johnston KR, et al. *Am J Med*. 1994; 97(2): 185–91.
16. Scientific Advisory Committee on Nutrition. *SACN Annual Report 2015*. Available at: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/522438/ACN_annual_report_2015.pdf. Accessed March 2018.
17. Kamarul Zaman M, et al. *World J Gastroenterol*. 2015; 21(17): 5372–81.
18. Elia M, et al. *Aliment Pharmacol Ther*. 2008; 27(2): 120–45.
19. Whelan K, Schneider S. *Curr Opin Gastroenterol*. 2011; 27(2): 152–9.
20. Slavin JL, Greenberg NA. *Nutrition*. 2003; 19(6): 549–52.
21. Tarleton SM, et al. *Practical Gastroenterology*. 2013; 11–22.
22. Halmos EP, et al. *Aliment Pharmacol Ther*. 2010; 32(7): 925–33.
23. Yoon SR, et al. *Nutr J*. 2015; 14: 116.
24. McClave SA, et al. *JPEN J Parenter Enteral Nutr*. 2016; 40(2): 159–211.



IDENTIFYING SYMPTOMS





IDENTIFYING SYMPTOMS



**NAUSEA/
VOMITING**

⊖



**EARLY
SATIETY**

⊕



REFLUX

⊕



**EPIGASTRIC
PAIN**

⊕



**ABDOMINAL PAIN/
BLOATING**

⊕



CONSTIPATION

⊕



DIARRHOEA

⊕

PATIENT HISTORY



POTENTIAL CAUSES

NUTRITIONAL CONSIDERATIONS

MEDICATION OPTIONS



IDENTIFYING SYMPTOMS



NAUSEA/
VOMITING

⊖



EARLY
SATIETY

⊕



REFLUX

⊕



EPIGASTRIC
PAIN

⊕



ABDOMINAL PAIN/
BLOATING

⊕



CONSTIPATION

⊕



DIARRHOEA

⊕

PATIENT HISTORY

Initiate 'vomit diary'

Rule out constipation (abdominal exam/X-ray/digital rectal examination).
If constipation present, follow *Constipation* section

POTENTIAL CAUSES

NUTRITIONAL CONSIDERATIONS

MEDICATION OPTIONS

IDENTIFYING SYMPTOMS



**NAUSEA/
VOMITING**

⊖



**EARLY
SATIETY**

⊕



REFLUX

⊕



**EPIGASTRIC
PAIN**

⊕



**ABDOMINAL PAIN/
BLOATING**

⊕



CONSTIPATION

⊕



DIARRHOEA

⊕

PATIENT HISTORY ⊕

POTENTIAL CAUSES ⊖

- **For vomiting:**
 - Rule out systemic infection or sepsis
 - Consult vomiting diary to identify triggers, e.g. hoisting, suctioning of secretions, tracheostomy care
 - Seek medical advice to continue or discontinue feeding
- Ensure feeding tube in correct position (nasojejunal tubes or jejunal extensions may migrate)
- Consider patient posture
- Consider whether symptoms occur with feeding or medications
- Establish if volume of fluid with medications is excessive
- Consider whether delayed gastric emptying may be causing symptoms

NUTRITIONAL CONSIDERATIONS ⊕

MEDICATION OPTIONS



IDENTIFYING SYMPTOMS



NAUSEA/
VOMITING



EARLY
SATIETY



REFLUX



EPIGASTRIC
PAIN



ABDOMINAL PAIN/
BLOATING



CONSTIPATION



DIARRHOEA



PATIENT HISTORY



POTENTIAL CAUSES



NUTRITIONAL CONSIDERATIONS



1st-line approaches:

- Reduce infusion rate of feed^{1,10}
- Change feed timing to avoid symptom triggers
- Consider bolus vs. continuous feeding
- Reduce feed volume¹⁰
- Reduce size of water flushes¹⁰
- Consider the side effects of medications, e.g. anticholinergics or anti-parkinsonians
- Ensure patient is elevated to ≥ 45 degrees

- Ensure feed timings do not reduce efficacy of medications, such as PPIs

2nd-line approaches:

- Consider trial of whey-based peptide formula to enhance gastric emptying^{4,5}
- Consider soy formula where intolerances are suspected

3rd-line approaches:

- Consider jejunal feeding¹

MEDICATION OPTIONS



PPI = proton pump inhibitor



IDENTIFYING SYMPTOMS



**NAUSEA/
VOMITING**

⊖



**EARLY
SATIETY**

⊕



REFLUX

⊕



**EPIGASTRIC
PAIN**

⊕



**ABDOMINAL PAIN/
BLOATING**

⊕



CONSTIPATION

⊕



DIARRHOEA

⊕

PATIENT HISTORY ⊕

POTENTIAL CAUSES ⊕

NUTRITIONAL CONSIDERATIONS ⊕

MEDICATION OPTIONS ⊖

Consider anti-emetic:

- Centrally acting antiemetic, e.g. cyclizine
- If related to movement, consider a peripheral acting anti-emetic, e.g. Stemetil (prochlorperazine)

References

1. Wang K, et al. *JPEN J Parenter Enteral Nutr.* 2017; 41(6): 959–67.
2. Whelan K, et al. *Eur J Clin Nutr.* 2004; 58(7): 1030–7.
3. King's College London. King's Stool Chart. Available at: www.kcl.ac.uk/stoolchart. Accessed March 2018.
4. Romano C, et al. *J Pediatr Gastroenterol Nutr.* 2017; 65(2): 242–264.
5. Alexander DD, et al. *World J Gastrointest Pharmacol Ther.* 2016; 7(2): 306–19.
6. Lee-Robichaud H, et al. *Cochrane Database Syst Rev.* 2010; 7: CD007570.
7. Minor G, et al. *Glob Pediatr Health.* 2016; 21;3:2333794X16681887.
8. Scientific Advisory Committee on Nutrition. *SACN Annual Report 2015.* Available at: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/522438/ACN_annual_report_2015.pdf. Accessed March 2018.
9. Todorovic V, Micklewright A (eds.) *A Pocket Guide to Clinical Nutrition.* Fourth Edition. The Parenteral and Enteral Nutrition Group: A Specialist Group of the British Dietetic Association, 2011.
10. Dietitians Association of Australia. *Enteral Nutrition Manual for Adults in Health Care Facilities.* DAA Nutrition Support Interest Group, 2015.
11. Barrett JS, et al. *JPEN J Parenter Enteral Nutr.* 2009; 33(1): 21–6.
12. National Institute for Health and Care Excellence. *Faecal Incontinence in Adults: Management.* NICE, 2007. Available at: <https://www.nice.org.uk/guidance/cg49>. Accessed March 2018.



IDENTIFYING SYMPTOMS

**NAUSEA/
VOMITING**
⊖

**EARLY
SATIETY**
⊕

REFLUX
⊕

**EPIGASTRIC
PAIN**
⊕

**ABDOMINAL PAIN/
BLOATING**
⊕

CONSTIPATION
⊕

DIARRHOEA
⊕

PATIENT HISTORY



POTENTIAL CAUSES



NUTRITIONAL CONSIDERATIONS

MEDICATION OPTIONS



IDENTIFYING SYMPTOMS



**NAUSEA/
VOMITING**

⊖



**EARLY
SATIETY**

⊕



REFLUX

⊕



**EPIGASTRIC
PAIN**

⊕



**ABDOMINAL PAIN/
BLOATING**

⊕



CONSTIPATION

⊕



DIARRHOEA

⊕

PATIENT HISTORY ⊕

POTENTIAL CAUSES ⊕

NUTRITIONAL CONSIDERATIONS ⊕

MEDICATION OPTIONS



IDENTIFYING SYMPTOMS

**NAUSEA/
VOMITING**

⊖

**EARLY
SATIETY**

⊕

REFLUX

⊕

**EPIGASTRIC
PAIN**

⊕

**ABDOMINAL PAIN/
BLOATING**

⊕

CONSTIPATION

⊕

DIARRHOEA

⊕

PATIENT HISTORY ⊕

POTENTIAL CAUSES ⊕

NUTRITIONAL CONSIDERATIONS ⊕

MEDICATION OPTIONS ⊕



IDENTIFYING SYMPTOMS

NAUSEA/
VOMITING
+

**EARLY
SATIETY**
-

REFLUX
+

EPIGASTRIC
PAIN
+

ABDOMINAL PAIN/
BLOATING
+

CONSTIPATION
+

DIARRHOEA
+

PATIENT HISTORY



POTENTIAL CAUSES

NUTRITIONAL CONSIDERATIONS

MEDICATION OPTIONS





IDENTIFYING SYMPTOMS



PATIENT HISTORY



Rule out constipation (abdominal exam/X-ray/digital rectal examination).
If constipation present, follow *Constipation* section

POTENTIAL CAUSES



NUTRITIONAL CONSIDERATIONS

MEDICATION OPTIONS



IDENTIFYING SYMPTOMS



PATIENT HISTORY (+)

POTENTIAL CAUSES (-)

- **For vomiting:**
 - Rule out systemic infection or sepsis
 - Consult vomiting diary to identify triggers, e.g. hoisting, suctioning of secretions, tracheostomy care
 - Seek medical advice to continue or discontinue feeding
- Ensure feeding tube in correct position (nasojejunal tubes or jejunal extensions may migrate)
- Consider patient posture
- Consider whether symptoms occur with feeding or medications
- Establish if volume of fluid with medications is excessive
- Consider whether delayed gastric emptying may be causing symptoms

NUTRITIONAL CONSIDERATIONS (+)

MEDICATION OPTIONS

IDENTIFYING SYMPTOMS



- PATIENT HISTORY (+)
- POTENTIAL CAUSES (+)
- NUTRITIONAL CONSIDERATIONS (-)**

1st-line approaches:

- Reduce infusion rate of feed^{1,10}
- Change feed timing to avoid symptom triggers
- Consider bolus vs. continuous feeding
- Reduce feed volume¹⁰
- Reduce size of water flushes¹⁰
- Consider the side effects of medications, e.g. anticholinergics or anti-parkinsonians
- Ensure patient is elevated to ≥ 45 degrees

- Ensure feed timings do not reduce efficacy of medications, such as PPIs

2nd-line approaches:

- Consider trial of whey-based peptide formula to enhance gastric emptying^{4,5}
- Consider soy formula where intolerances are suspected

3rd-line approaches:

- Consider jejunal feeding¹

- MEDICATION OPTIONS (+)

PPI = proton pump inhibitor

IDENTIFYING SYMPTOMS



- PATIENT HISTORY (+)
- POTENTIAL CAUSES (+)
- NUTRITIONAL CONSIDERATIONS (+)
- MEDICATION OPTIONS (-)**

Consider prokinetic,
 e.g. erythromycin (avoid metoclopramide in Parkinson’s disease and traumatic brain injury)



IDENTIFYING SYMPTOMS



NAUSEA/
VOMITING

+



EARLY
SATIETY

-



REFLUX

+



EPIGASTRIC
PAIN

+



ABDOMINAL PAIN/
BLOATING

+



CONSTIPATION

+



DIARRHOEA

+

PATIENT HISTORY



POTENTIAL CAUSES



NUTRITIONAL CONSIDERATIONS

MEDICATION OPTIONS





IDENTIFYING SYMPTOMS



NAUSEA/
VOMITING

+



EARLY
SATIETY

-



REFLUX

+



EPIGASTRIC
PAIN

+



ABDOMINAL PAIN/
BLOATING

+



CONSTIPATION

+



DIARRHOEA

+

PATIENT HISTORY +

POTENTIAL CAUSES +

NUTRITIONAL CONSIDERATIONS +

MEDICATION OPTIONS



IDENTIFYING SYMPTOMS

NAUSEA/
VOMITING

+

EARLY
SATIETY

-

REFLUX

+

EPIGASTRIC
PAIN

+

ABDOMINAL PAIN/
BLOATING

+

CONSTIPATION

+

DIARRHOEA

+

PATIENT HISTORY +

POTENTIAL CAUSES +

NUTRITIONAL CONSIDERATIONS +

MEDICATION OPTIONS +



IDENTIFYING SYMPTOMS



NAUSEA/
VOMITING

+



EARLY
SATIETY

+



REFLUX

-



EPIGASTRIC
PAIN

+



ABDOMINAL PAIN/
BLOATING

+



CONSTIPATION

+



DIARRHOEA

+

PATIENT HISTORY +

POTENTIAL CAUSES

NUTRITIONAL CONSIDERATIONS

MEDICATION OPTIONS





IDENTIFYING SYMPTOMS



PATIENT HISTORY



Rule out constipation (abdominal exam/X-ray/digital rectal examination).
If constipation present, follow *Constipation* section

POTENTIAL CAUSES



NUTRITIONAL CONSIDERATIONS

MEDICATION OPTIONS



IDENTIFYING SYMPTOMS



- PATIENT HISTORY (+)
- POTENTIAL CAUSES (-)**

- **For vomiting:**
 - Rule out systemic infection or sepsis
 - Consult vomiting diary to identify triggers, e.g. hoisting, suctioning of secretions, tracheostomy care
 - Seek medical advice to continue or discontinue feeding
- Ensure feeding tube in correct position (nasojejunal tubes or jejunal extensions may migrate)
- Consider patient posture
- Consider whether symptoms occur with feeding or medications
- Establish if volume of fluid with medications is excessive
- Consider whether delayed gastric emptying may be causing symptoms

- NUTRITIONAL CONSIDERATIONS (+)
- MEDICATION OPTIONS



IDENTIFYING SYMPTOMS



- PATIENT HISTORY (+)
- POTENTIAL CAUSES (+)
- NUTRITIONAL CONSIDERATIONS (-)**

1st-line approaches:

- Reduce infusion rate of feed^{1,10}
- Change feed timing to avoid symptom triggers
- Consider bolus vs. continuous feeding
- Reduce feed volume¹⁰
- Reduce size of water flushes¹⁰
- Consider the side effects of medications, e.g. anticholinergics or anti-parkinsonians
- Ensure patient is elevated to ≥ 45 degrees

- Ensure feed timings do not reduce efficacy of medications, such as PPIs

2nd-line approaches:

- Consider trial of whey-based peptide formula to enhance gastric emptying^{4,5}
- Consider soy formula where intolerances are suspected

3rd-line approaches:

- Consider jejunal feeding¹

- MEDICATION OPTIONS (+)

PPI = proton pump inhibitor

IDENTIFYING SYMPTOMS



- PATIENT HISTORY (+)
- POTENTIAL CAUSES (+)
- NUTRITIONAL CONSIDERATIONS (+)
- MEDICATION OPTIONS (-)**

- **Optimise PPI dose:**
 - Switch PPI
 - Double dose
- Add H2 antagonist at night, e.g. ranitidine
- Antacid (alginate) at start/end of feed
- Consider prokinetic (avoid metoclopramide in Parkinson's disease and traumatic brain injury)



IDENTIFYING SYMPTOMS



NAUSEA/
VOMITING

+



EARLY
SATIETY

+



REFLUX

-



EPIGASTRIC
PAIN

+



ABDOMINAL PAIN/
BLOATING

+



CONSTIPATION

+



DIARRHOEA

+

PATIENT HISTORY +

POTENTIAL CAUSES +

NUTRITIONAL CONSIDERATIONS

MEDICATION OPTIONS



IDENTIFYING SYMPTOMS



NAUSEA/
VOMITING

+



EARLY
SATIETY

+



REFLUX

-



EPIGASTRIC
PAIN

+



ABDOMINAL PAIN/
BLOATING

+



CONSTIPATION

+



DIARRHOEA

+

PATIENT HISTORY +

POTENTIAL CAUSES +

NUTRITIONAL CONSIDERATIONS +

MEDICATION OPTIONS





IDENTIFYING SYMPTOMS

NAUSEA/
VOMITING

+

EARLY
SATIETY

+

REFLUX

-

EPIGASTRIC
PAIN

+

ABDOMINAL PAIN/
BLOATING

+

CONSTIPATION

+

DIARRHOEA

+

PATIENT HISTORY +

POTENTIAL CAUSES +

NUTRITIONAL CONSIDERATIONS +

MEDICATION OPTIONS +





IDENTIFYING SYMPTOMS

NAUSEA/
VOMITING

+

EARLY
SATIETY

+

REFLUX

+

EPIGASTRIC
PAIN

-

ABDOMINAL PAIN/
BLOATING

+

CONSTIPATION

+

DIARRHOEA

+

PATIENT HISTORY +

POTENTIAL CAUSES

NUTRITIONAL CONSIDERATIONS

MEDICATION OPTIONS



IDENTIFYING SYMPTOMS

NAUSEA/
VOMITING

+

EARLY
SATIETY

+

REFLUX

+

EPIGASTRIC
PAIN

-

ABDOMINAL PAIN/
BLOATING

+

CONSTIPATION

+

DIARRHOEA

+

PATIENT HISTORY -

Rule out constipation (abdominal exam/X-ray/digital rectal examination)

POTENTIAL CAUSES +

NUTRITIONAL CONSIDERATIONS

MEDICATION OPTIONS



IDENTIFYING SYMPTOMS



PATIENT HISTORY



POTENTIAL CAUSES



NO CONSTIPATION



CONSTIPATION PRESENT



NUTRITIONAL CONSIDERATIONS

MEDICATION OPTIONS





IDENTIFYING SYMPTOMS



PATIENT HISTORY (+)

POTENTIAL CAUSES (-)

NO CONSTIPATION (-)

Consider:

- Functional causes, e.g. gastric emptying delay, chronic intestinal pseudo-obstruction, megacolon
- Organic causes, e.g. small intestinal bacterial overgrowth, gallstones, hiatus hernia, peptic ulcer disease, pancreatitis, fatty liver

CONSTIPATION PRESENT (→)

NUTRITIONAL CONSIDERATIONS (+)

MEDICATION OPTIONS



IDENTIFYING SYMPTOMS



- PATIENT HISTORY (+)
- POTENTIAL CAUSES (+)
- NUTRITIONAL CONSIDERATIONS (-)**

1st-line approaches:

- Reduce infusion rate of feed^{1,10}
- Change feed timing to avoid symptom triggers
- Consider bolus vs. continuous feeding
- Reduce feed volume¹⁰
- Reduce size of water flushes¹⁰
- Consider the side effects of medications, e.g. anticholinergics or anti-parkinsonians
- Ensure patient is elevated to ≥ 45 degrees

- Ensure feed timings do not reduce efficacy of medications, such as PPIs

2nd-line approaches:

- Consider trial of whey-based peptide formula to enhance gastric emptying^{4,5}
- Consider soy formula where intolerances are suspected

3rd-line approaches:

- Consider jejunal feeding¹

- MEDICATION OPTIONS (+)

PPI = proton pump inhibitor

IDENTIFYING SYMPTOMS



- PATIENT HISTORY (+)
- POTENTIAL CAUSES (+)
- NUTRITIONAL CONSIDERATIONS (+)
- MEDICATION OPTIONS (-)**

- **Optimise PPI dose:**
 - Switch PPI
 - Double dose
- Add H2 antagonist at night, e.g. ranitidine
- Antacid (alginate) at start/end of feed
- Consider prokinetic (avoid metoclopramide in Parkinson's disease and traumatic brain injury)
- Stop lactulose⁶
- Consider low-dose tricyclic



IDENTIFYING SYMPTOMS

NAUSEA/
VOMITING

+

EARLY
SATIETY

+

REFLUX

+

EPIGASTRIC
PAIN

-

ABDOMINAL PAIN/
BLOATING

+

CONSTIPATION

+

DIARRHOEA

+

PATIENT HISTORY +

POTENTIAL CAUSES +

NUTRITIONAL CONSIDERATIONS

MEDICATION OPTIONS



IDENTIFYING SYMPTOMS

NAUSEA/
VOMITING

+

EARLY
SATIETY

+

REFLUX

+

EPIGASTRIC
PAIN

-

ABDOMINAL PAIN/
BLOATING

+

CONSTIPATION

+

DIARRHOEA

+

PATIENT HISTORY



POTENTIAL CAUSES



NUTRITIONAL CONSIDERATIONS



MEDICATION OPTIONS





IDENTIFYING SYMPTOMS



NAUSEA/
VOMITING

+



EARLY
SATIETY

+



REFLUX

+



EPIGASTRIC
PAIN

-



ABDOMINAL PAIN/
BLOATING

+



CONSTIPATION

+



DIARRHOEA

+

PATIENT HISTORY



POTENTIAL CAUSES



NUTRITIONAL CONSIDERATIONS



MEDICATION OPTIONS





IDENTIFYING SYMPTOMS



NAUSEA/
VOMITING

+



EARLY
SATIETY

+



REFLUX

+



EPIGASTRIC
PAIN

+



ABDOMINAL PAIN/
BLOATING

-



CONSTIPATION

+



DIARRHOEA

+

PATIENT HISTORY



POTENTIAL CAUSES

NUTRITIONAL CONSIDERATIONS

MEDICATION OPTIONS



IDENTIFYING SYMPTOMS



PATIENT HISTORY -

Rule out constipation (abdominal exam/X-ray/digital rectal examination).

POTENTIAL CAUSES +

NUTRITIONAL CONSIDERATIONS

MEDICATION OPTIONS



IDENTIFYING SYMPTOMS



NAUSEA/
VOMITING

+



EARLY
SATIETY

+



REFLUX

+



EPIGASTRIC
PAIN

+



ABDOMINAL PAIN/
BLOATING

-



CONSTIPATION

+



DIARRHOEA

+

- PATIENT HISTORY +
- POTENTIAL CAUSES** -
- NO CONSTIPATION +
- CONSTIPATION PRESENT →
- NUTRITIONAL CONSIDERATIONS
- MEDICATION OPTIONS





IDENTIFYING SYMPTOMS



PATIENT HISTORY



POTENTIAL CAUSES



NO CONSTIPATION



Consider:

- Functional causes, e.g. gastric emptying delay, chronic intestinal pseudo-obstruction, megacolon
- Organic causes, e.g. small intestinal bacterial overgrowth, gallstones, hiatus hernia, peptic ulcer disease, pancreatitis, fatty liver

CONSTIPATION PRESENT



NUTRITIONAL CONSIDERATIONS



MEDICATION OPTIONS



IDENTIFYING SYMPTOMS



- PATIENT HISTORY (+)
- POTENTIAL CAUSES (+)
- NUTRITIONAL CONSIDERATIONS (-)**

- Trial fibre-containing vs. fibre-free formulas⁹
- Consider additional fibre supplements if required (optimal fibre intake for a healthy adult is 25–30g/day)⁸
- Consider whey-based peptide feed if pre-morbid/persistent constipation⁷
- Be aware of the type of fibre used; FODMAP-containing formulas may worsen symptoms of bloating or diarrhoea in sensitive patients¹¹

- MEDICATION OPTIONS (+)

IDENTIFYING SYMPTOMS



- PATIENT HISTORY (+)
- POTENTIAL CAUSES (+)
- NUTRITIONAL CONSIDERATIONS (+)
- MEDICATION OPTIONS (-)**

CONSTIPATION

- **Rectal evacuation difficulties:**
 - Suppositories: glycerin or bisacodyl or leccarbon or Microlax® (order of medications based on escalating levels of stimulation of rectal evacuation)
- **Slow transit:**
 - Senna/softener
 - Macrogol (avoid lactulose)
 - Prokinetics
- **Stop lactulose⁶**
- **For epigastric pain,** consider low-dose tricyclic
- **For abdominal pain and bloating,** consider venting of air from gastrostomy tube (if in situ)



IDENTIFYING SYMPTOMS



NAUSEA/
VOMITING

+



EARLY
SATIETY

+



REFLUX

+



EPIGASTRIC
PAIN

+



ABDOMINAL PAIN/
BLOATING

-



CONSTIPATION

+



DIARRHOEA

+

PATIENT HISTORY +

POTENTIAL CAUSES +

NUTRITIONAL CONSIDERATIONS

MEDICATION OPTIONS





IDENTIFYING SYMPTOMS



NAUSEA/
VOMITING

+



EARLY
SATIETY

+



REFLUX

+



EPIGASTRIC
PAIN

+



ABDOMINAL PAIN/
BLOATING

-



CONSTIPATION

+



DIARRHOEA

+

- PATIENT HISTORY +
- POTENTIAL CAUSES +
- NUTRITIONAL CONSIDERATIONS +
- MEDICATION OPTIONS





IDENTIFYING SYMPTOMS



NAUSEA/
VOMITING

+



EARLY
SATIETY

+



REFLUX

+



EPIGASTRIC
PAIN

+



ABDOMINAL PAIN/
BLOATING

-



CONSTIPATION

+



DIARRHOEA

+

- PATIENT HISTORY +
- POTENTIAL CAUSES +
- NUTRITIONAL CONSIDERATIONS +
- MEDICATION OPTIONS +



IDENTIFYING SYMPTOMS

NAUSEA/
VOMITING

+

EARLY
SATIETY

+

REFLUX

+

EPIGASTRIC
PAIN

+

ABDOMINAL PAIN/
BLOATING

+

CONSTIPATION

-

DIARRHOEA

+

PATIENT HISTORY



POTENTIAL CAUSES

NUTRITIONAL CONSIDERATIONS

MEDICATION OPTIONS



IDENTIFYING SYMPTOMS



PATIENT HISTORY

Patient history; consult King's Stool Chart^{2,3} (or Bristol Stool Chart)

POTENTIAL CAUSES

NUTRITIONAL CONSIDERATIONS

MEDICATION OPTIONS

IDENTIFYING SYMPTOMS



PATIENT HISTORY 

POTENTIAL CAUSES 

Rule out:

- Rectal evacuation difficulties
- Slow transit
- Both

- Consider posture when toileting and psychosocial issues
- Ensure adequate hydration and consider 'free' water content of enteral formula

NUTRITIONAL CONSIDERATIONS 

MEDICATION OPTIONS

IDENTIFYING SYMPTOMS



PATIENT HISTORY 

POTENTIAL CAUSES 

NUTRITIONAL CONSIDERATIONS 

- Trial fibre-containing vs. fibre-free formulas⁹
- Consider additional fibre supplements if required (optimal fibre intake for a healthy adult is 25–30g/day)⁸
- Consider whey-based peptide feed if pre-morbid/persistent constipation⁷
- Be aware of the type of fibre used; FODMAP-containing formulas may worsen symptoms of bloating or diarrhoea in sensitive patients¹¹

MEDICATION OPTIONS 

IDENTIFYING SYMPTOMS



- PATIENT HISTORY (+)
- POTENTIAL CAUSES (+)
- NUTRITIONAL CONSIDERATIONS (+)
- MEDICATION OPTIONS (-)**

CONSTIPATION

- **Rectal evacuation difficulties:**
 - Suppositories: glycerin or bisacodyl or leccarbon or Microlax® (order of medications based on escalating levels of stimulation of rectal evacuation)
- **Slow transit:**
 - Senna/softener
 - Macrogol (avoid lactulose)
 - Prokinetics

IDENTIFYING SYMPTOMS



NAUSEA/
VOMITING

+



EARLY
SATIETY

+



REFLUX

+



EPIGASTRIC
PAIN

+



ABDOMINAL PAIN/
BLOATING

+



CONSTIPATION

-



DIARRHOEA

+

PATIENT HISTORY 

POTENTIAL CAUSES 

NUTRITIONAL CONSIDERATIONS

MEDICATION OPTIONS



IDENTIFYING SYMPTOMS



NAUSEA/
VOMITING

+



EARLY
SATIETY

+



REFLUX

+



EPIGASTRIC
PAIN

+



ABDOMINAL PAIN/
BLOATING

+



CONSTIPATION

-



DIARRHOEA

+

PATIENT HISTORY +

POTENTIAL CAUSES +

NUTRITIONAL CONSIDERATIONS +

MEDICATION OPTIONS



IDENTIFYING SYMPTOMS



NAUSEA/
VOMITING

+



EARLY
SATIETY

+



REFLUX

+



EPIGASTRIC
PAIN

+



ABDOMINAL PAIN/
BLOATING

+



CONSTIPATION

-



DIARRHOEA

+

PATIENT HISTORY +

POTENTIAL CAUSES +

NUTRITIONAL CONSIDERATIONS +

MEDICATION OPTIONS +



IDENTIFYING SYMPTOMS



NAUSEA/
VOMITING

+



EARLY
SATIETY

+



REFLUX

+



EPIGASTRIC
PAIN

+



ABDOMINAL PAIN/
BLOATING

+



CONSTIPATION

+



DIARRHOEA

-

PATIENT HISTORY



POTENTIAL CAUSES

NUTRITIONAL CONSIDERATIONS

MEDICATION OPTIONS





IDENTIFYING SYMPTOMS



PATIENT HISTORY



Patient history; consult King's Stool Chart^{2,3} (or Bristol Stool Chart)

POTENTIAL CAUSES



NUTRITIONAL CONSIDERATIONS

MEDICATION OPTIONS





IDENTIFYING SYMPTOMS



PATIENT HISTORY



POTENTIAL CAUSES



Rule out:

- Infection/overflow diarrhoea⁹
- Medication, e.g. antibiotics, sorbitol-containing liquids
- Organic, e.g. IBD
- Functional, e.g. IBS
- Faecal incontinence
- Rectal prolapse (Acute cases require referral)

IBD = inflammatory bowel disease; IBS = irritable bowel syndrome

NUTRITIONAL CONSIDERATIONS



MEDICATION OPTIONS



IDENTIFYING SYMPTOMS



- PATIENT HISTORY +
- POTENTIAL CAUSES +
- NUTRITIONAL CONSIDERATIONS** -

- Trial fibre-containing vs. fibre-free formulas⁹
- Consider additional fibre supplements if required (optimal fibre intake for a healthy adult is 25–30g/day)⁸
- Consider whey-based peptide feed if pre-morbid/persistent constipation⁷
- Be aware of the type of fibre used; FODMAP-containing formulas may worsen symptoms of bloating or diarrhoea in sensitive patients¹¹
- Trial reduction in feed rate or time on feed (e.g. using a lower volume formula)
- Trial peptide feed and/or lower osmolarity formula for diarrhoea^{9,11}

FODMAP = fermentable oligosaccharides, disaccharides, monosaccharides and polyols

MEDICATION OPTIONS +

IDENTIFYING SYMPTOMS



- PATIENT HISTORY (+)
- POTENTIAL CAUSES (+)
- NUTRITIONAL CONSIDERATIONS (+)
- MEDICATION OPTIONS (-)**

DIARRHOEA

- Add/optimize anti-diarrhoeal
- Overflow:
 - Suppository or enema
 - Oral Macrogol
 - Optimize medication
- Faecal incontinence:
 - Follow NICE CG49¹²
 - Optimize medication



IDENTIFYING SYMPTOMS



NAUSEA/
VOMITING

+



EARLY
SATIETY

+



REFLUX

+



EPIGASTRIC
PAIN

+



ABDOMINAL PAIN/
BLOATING

+



CONSTIPATION

+



DIARRHOEA

-

PATIENT HISTORY



POTENTIAL CAUSES



NUTRITIONAL CONSIDERATIONS

MEDICATION OPTIONS





IDENTIFYING SYMPTOMS



NAUSEA/
VOMITING

+



EARLY
SATIETY

+



REFLUX

+



EPIGASTRIC
PAIN

+



ABDOMINAL PAIN/
BLOATING

+



CONSTIPATION

+



DIARRHOEA

-

PATIENT HISTORY



POTENTIAL CAUSES



NUTRITIONAL CONSIDERATIONS



MEDICATION OPTIONS





IDENTIFYING SYMPTOMS



NAUSEA/
VOMITING

+



EARLY
SATIETY

+



REFLUX

+



EPIGASTRIC
PAIN

+



ABDOMINAL PAIN/
BLOATING

+



CONSTIPATION

+



DIARRHOEA

-

PATIENT HISTORY



POTENTIAL CAUSES



NUTRITIONAL CONSIDERATIONS



MEDICATION OPTIONS

